


<b>TRIUMF - RESEARCH PROPOSAL</b> 		<b>Experiment no</b> <b>E1073</b>	<b>Sheet 1 of 15</b>																																																																																
<b>Title of proposed experiment</b> Ground state mass determination of the halo nucleus $^{11}\text{Li}$																																																																																			
<b>Name of group</b> TITAN																																																																																			
<b>Spokesperson for group</b> Jens Dilling and Fred Sarazin																																																																																			
<b>Email address</b> JDilling@triumf.ca and FSarazin@mines.edu																																																																																			
<table border="0"> <tr> <td>J. Dilling</td> <td>TRIUMF</td> <td>Scientist</td> <td>80 %</td> </tr> <tr> <td>F. Sarazin</td> <td>Colorado School of Mines</td> <td>Professor</td> <td>30 %</td> </tr> <tr> <td>P. Delheij</td> <td>TRIUMF</td> <td>Scientist</td> <td>80 %</td> </tr> <tr> <td>P. Bricault</td> <td>TRIUMF</td> <td>Scientist</td> <td>20 %</td> </tr> <tr> <td>J. Caggiano</td> <td>TRIUMF</td> <td>Scientist</td> <td>30 %</td> </tr> <tr> <td>J. Lee</td> <td>McGill</td> <td>Professor</td> <td>30%</td> </tr> <tr> <td>J. Crawford</td> <td>McGill</td> <td>Professor</td> <td>30%</td> </tr> <tr> <td>F. Buchinger</td> <td>McGill</td> <td>Lecturer</td> <td>30%</td> </tr> <tr> <td>B. Moore</td> <td>McGill</td> <td>Professor</td> <td>30%</td> </tr> <tr> <td>G. Gwinner</td> <td>U. of Manitoba</td> <td>Professor</td> <td>40%</td> </tr> <tr> <td>M. Pearson</td> <td>TRIUMF</td> <td>Scientist</td> <td>30 %</td> </tr> <tr> <td>M. Smith</td> <td>UBC</td> <td>Ph.D. Student</td> <td>80 %</td> </tr> <tr> <td>Z. Ke</td> <td>U. of Manitoba</td> <td>Ph.D. Student</td> <td>80 %</td> </tr> <tr> <td>M. Brodeur</td> <td>UBC</td> <td>M.Sc. Student</td> <td>80%</td> </tr> <tr> <td>L. Bromley</td> <td>McGill University</td> <td>M.Sc. Student</td> <td>80%</td> </tr> <tr> <td>Caleb Mattoon</td> <td>Colorado School of Mines</td> <td>Ph.D. Student</td> <td>20%</td> </tr> <tr> <td>Matt Porten-Peden</td> <td>Colorado School of Mines</td> <td>Ph.D. Student</td> <td>30%</td> </tr> <tr> <td>J. Vaz</td> <td>TRIUMF</td> <td>Post-doc</td> <td>100%</td> </tr> <tr> <td>V. Ryjkov</td> <td>TRIUMF</td> <td>Post-doc</td> <td>100%</td> </tr> <tr> <td>G. Sikler</td> <td>TRIUMF</td> <td>Post-doc</td> <td>100%</td> </tr> </table>				J. Dilling	TRIUMF	Scientist	80 %	F. Sarazin	Colorado School of Mines	Professor	30 %	P. Delheij	TRIUMF	Scientist	80 %	P. Bricault	TRIUMF	Scientist	20 %	J. Caggiano	TRIUMF	Scientist	30 %	J. Lee	McGill	Professor	30%	J. Crawford	McGill	Professor	30%	F. Buchinger	McGill	Lecturer	30%	B. Moore	McGill	Professor	30%	G. Gwinner	U. of Manitoba	Professor	40%	M. Pearson	TRIUMF	Scientist	30 %	M. Smith	UBC	Ph.D. Student	80 %	Z. Ke	U. of Manitoba	Ph.D. Student	80 %	M. Brodeur	UBC	M.Sc. Student	80%	L. Bromley	McGill University	M.Sc. Student	80%	Caleb Mattoon	Colorado School of Mines	Ph.D. Student	20%	Matt Porten-Peden	Colorado School of Mines	Ph.D. Student	30%	J. Vaz	TRIUMF	Post-doc	100%	V. Ryjkov	TRIUMF	Post-doc	100%	G. Sikler	TRIUMF	Post-doc	100%
J. Dilling	TRIUMF	Scientist	80 %																																																																																
F. Sarazin	Colorado School of Mines	Professor	30 %																																																																																
P. Delheij	TRIUMF	Scientist	80 %																																																																																
P. Bricault	TRIUMF	Scientist	20 %																																																																																
J. Caggiano	TRIUMF	Scientist	30 %																																																																																
J. Lee	McGill	Professor	30%																																																																																
J. Crawford	McGill	Professor	30%																																																																																
F. Buchinger	McGill	Lecturer	30%																																																																																
B. Moore	McGill	Professor	30%																																																																																
G. Gwinner	U. of Manitoba	Professor	40%																																																																																
M. Pearson	TRIUMF	Scientist	30 %																																																																																
M. Smith	UBC	Ph.D. Student	80 %																																																																																
Z. Ke	U. of Manitoba	Ph.D. Student	80 %																																																																																
M. Brodeur	UBC	M.Sc. Student	80%																																																																																
L. Bromley	McGill University	M.Sc. Student	80%																																																																																
Caleb Mattoon	Colorado School of Mines	Ph.D. Student	20%																																																																																
Matt Porten-Peden	Colorado School of Mines	Ph.D. Student	30%																																																																																
J. Vaz	TRIUMF	Post-doc	100%																																																																																
V. Ryjkov	TRIUMF	Post-doc	100%																																																																																
G. Sikler	TRIUMF	Post-doc	100%																																																																																
<b>Date for start of preparations:</b> Summer 2006		<b>Beam time requested:</b> <b>12-hr shifts</b> <b>Beam line/channel</b> <b>Polarized primary beam?</b>																																																																																	
<b>Date ready:</b> Aug. 2006		6 12h shifts of stable $^7\text{Li}$ at 30 keV from OLIS 4 12h shifts of $^{11}\text{Li}$ at 30 keV from ISAC																																																																																	

Completion date:

date

The halo occurrence observed in light neutron-rich nuclei, such as  $^{11}\text{Li}$ , is among the most exciting phenomena observed far away from stability. It is a remarkable behavior from the nuclear physics point-of-view, but more generally it is an example of quantum behavior in a microscopic few-body system, where some of its weakly bound components are venturing (tunneling) beyond the binding potential of the core. Since the first experiment pointing out the extended nature of the  $^{11}\text{Li}$  nucleus [TAN85], much work both experimentally and theoretically (see [JEN04] for a recent review on halo physics) have been performed to study and to understand the halo nature of  $^{11}\text{Li}$  (and other nuclei). The halo phenomenon was first suggested by Hansen and Jonson [HAN87]. In their simple model, they link the extended matter radius of  $^{11}\text{Li}$  to the weak binding energy of its last two neutrons ( $S_{2n}$ ). The matter radius decays as an exponential with a length equal to  $\hbar/(2\mu S_{2n})^{1/2}$ . Although more advanced models now exist, the binding energy of  $^{11}\text{Li}$  remains a key ingredient of those models and the calculated matter radius is still (very) sensitive to the two-neutron separation energy. With models of increasing complexity, it is highly desirable to measure the mass of  $^{11}\text{Li}$  within a few hundred eV.

**Experimental area**

TITAN  
ISAC I

**Primary beam and target** (energy, energy spread, intensity, pulse characteristics, emittance)

**Secondary channel**

Stable  ${}^7\text{Li}$  from OLIS at 30 keV of at least  $1 \cdot 10^5$  ions/s  
 ${}^{11}\text{Li}$  from ISAC at 30 keV of at least  $1 \cdot 10^4$  ions/s

**Secondary beam** (particle type, momentum range, momentum bite, solid angle, spot size, emittance, intensity, beam purity, target, special characteristics)

**TRIUMF SUPPORT:**

Summarize all equipment and technical support to be provided by TRIUMF. If new equipment is required, provide cost estimates.  
NOTE: Technical Review Forms must also be provided before allocation of beam time.

**NON-TRIUMF SUPPORT:**

Summarize the expected sources of funding for the experiment.

Identify major capital items and their costs that will be provided from these funds.

Summarize possible hazards associated with the experimental apparatus, precautions to be taken, and other matters that should be brought to the notice of the Safety Officer. Details must be provided separately in a safety report to be prepared by the spokesperson under the guidance of the Safety Report Guide available from the Science Division Office.

Taking the ISAC  $^{11}\text{Li}$  beam in continuous mode gives a neutron field around the beam dump (be this the RFQ or Penning trap) of  $10\mu\text{Sv}/\text{Hr}$  at 0.5m. This requires general flagging and ideally borated paraffin shielding, as used at E996 (GSI experiment).

General TITAN safety (HV).

(A) Scientific Motivation:

The halo occurrence observed in light neutron-rich nuclei, such as  $^{11}\text{Li}$ , is among the most exciting phenomena observed far away from stability. It is a remarkable behavior from the nuclear physics point-of-view, but more generally it is an example of quantum behavior in a microscopic few-body system, where some of its weakly bound components are venturing (tunneling) beyond the binding potential of the core. Since the first experiment pointing out the extended nature of the  $^{11}\text{Li}$  nucleus [TAN85], much work both experimentally and theoretically (see [JEN04] for a recent review on halo physics) have been performed to study and to understand the halo nature of  $^{11}\text{Li}$  (and other nuclei). The halo phenomenon was first suggested by Hansen and Jonson [HAN87]. In their simple model, they link the extended matter radius of  $^{11}\text{Li}$  to the weak binding energy of its last two neutrons ( $S_{2n}$ ). The matter radius decays as an exponential with a length equal to  $\hbar/(2\mu S_{2n})^{1/2}$ . Although more advanced models now exist, the binding energy of  $^{11}\text{Li}$  remains a key ingredient of those models and the calculated matter radius is still (very) sensitive to the two-neutron separation energy. With models of increasing complexity, it is highly desirable to measure the mass of  $^{11}\text{Li}$  within a few hundred eV.

Last year, the MISTRAL collaboration (based at ISOLDE – CERN) reported at the ENAM’04 conference [BAC04] the first precise mass measurement of  $^{11}\text{Li}$  from which they extracted the  $S_{2n}$  of the two halo neutrons.

While the MISTRAL measurement does not appear particularly anomalous, it is about 76 keV away from the 2003 Atomic Mass Evaluation (AME) [AUD03], which corresponds to a change of about 20% of the binding energy of the two halo neutrons. As the MISTRAL measurement is so much more precise than the other existing measurements, it is expected to pin down the mass and the  $S_{2n}$  for  $^{11}\text{Li}$  in the next mass compilation. A possible issue here is that the next generation of halo models may be based on a single precise experimental determination of the  $^{11}\text{Li}$  binding energy. And, in this context, it is worth noting that a preliminary and unpublished  $^{11}\text{Li}$  mass measurement at MISTRAL made in 2002 placed the mass of  $^{11}\text{Li}$  within the limits of AME’03 with a precision of  $\pm 25\text{keV}$  [LUN03].




Figure 1: Comparison of the various  $S_{2n}$  (mass) measurements for  $^{11}\text{Li}$  (from [BAC04])

This 2002 measurement is about  $3\sigma$  away from the present MISTRAL estimate. It therefore makes sense that a measurement with a similar precision using a different experimental setup should be undertaken to confirm the MISTRAL measurement. It is also important that this new measurement could be performed in a timely manner, as the binding energy of  $^{11}\text{Li}$  is a key ingredient in so many models, some of them based on new experimental results. For example, the charge radius of  $^{11}\text{Li}$  can be deduced from the total isotope shift for  $^{11}\text{Li}$ , which was recently measured at TRIUMF-ISAC [NOR05a]. The  $^{11}\text{Li}$  binding energy is here needed to calculate the mass dependent isotope shift, which is subtracted to the measured total isotope shift to give the  $^{11}\text{Li}$  charge radius. The effect of the new mass measurement by MISTRAL is quite dramatic (35% change) [NOR05b]. We propose to use the TITAN facility at ISAC/TRIUMF to measure the mass of  $^{11}\text{Li}$  by using the superior  $^{11}\text{Li}$  yield available at ISAC.

#### References:

- [AUD03] G.Audi et al., Nucl. Phys. A729 (2003) 337
- [BAC04] C.Bachelet, ENAM'04 conference; Eur. Phys. J. A25 (2005) 31
- [JEN04] A.S.Jensen, K.Riisager and D.V.Fedorov, Rev. Mod. Phys. 76 (2004) 215
- [HAN87] P.G.Hansen and B.Jonson, Europhys. Lett. 4 (1987) 409
- [LUN03] D.Lunney et al., Addendum to proposal IS-402 (2003) Unpublished
- [NOR05a] W.Nörtershauser et al., submitted to Phys. Rev. Lett. (2005) and physics/0509265
- [NOR05b] W.Nörtershauser, Private Communication (2005)
- [TAN85] I.Tanihata et al., Phys. Rev. Lett. 55 (1985) 2676 ; Phys. Lett. B160 (1985) 380

## (B,C) Description of the measurement technique and the experimental procedure

A detailed description of the TITAN facility and the experimental procedure is given in the EEC proposal "High accuracy mass measurements for superallowed nuclear  $\beta$ -decay emitters".

A schematic of the TITAN facility is shown in figure 2 below.

Legend:

ISO: ion source

RFCT: RF cooler trap

EBIT: electron beam ion trap

WIFI: Wien filter

CPET: cooler Penning trap

MPET: Measurement Penning trap




Fig. 2 Schematic of the TITAN setup.

The presently available production rate of  $^{11}\text{Li}$  available at ISAC-TRIUMF (about 44,000  $^{11}\text{Li}$  per second measured at the yield station) makes it possible to measure its binding energy, despite its 8.6ms half-life. For a mass determination in a single charge state, a relative accuracy of  $\delta m/m \approx 5 \cdot 10^{-8}$  at an excitation time of 20 ms and with a total cycle number of  $N=10\,000$ , could be reached. In this case, a single charge state would be sufficient, since only a factor of 3 is gained, and losses due to the breeding process can be avoided. Also additional filtering is not needed, since the Li beam is extremely clean. The absolute mass uncertainty would be 550 eV.

The procedure would include the bunching and cooling of Li in the TITAN RFQ, which is filled with helium gas. Since the He and Li masses are rather close, higher losses due to hard-sphere collisions are anticipated. These collisions scatter the charged Li ions either out of the RF-phase and lead to unstable trajectories, or the scattering leads to collisions with the RFQ rod structure directly.

Instead of the regular 50 % transmission for the RFCT, a conservative number of 1% is assumed. The remaining ions bypass the EBIT and go through WIFI2, where a moderate suppression of contamination can be achieved. Production of contamination could arise from charge exchange processes in the RFCT.

Moreover, since no charge-breeding is foreseen, the Li ions could be directly transferred into the measurement trap (MPET), hence the experiment could be done, even without the cooler Penning trap (The cooler trap will be brought to TRIUMF spring 2007. It is presently being designed, built and tested at the University of Manitoba).

(D) Readiness

The TITAN facility is presently under construction and first on-line measurements are planned for the second half of the year 2006. At that time the system will be without the Cooler Penning Trap (CPET). However, measurements of isotopes such as Li are possible and could be carried out at that stage.

The system and procedure could be tested in an ideal way, by providing Li beam from the off-line OLIS ion source and establish the transfer efficiency, including the cooling in the RFCT. Moreover the mass of  ${}^7\text{Li}$  is known to 80 eV or  $\delta m/m = 2 \cdot 10^{-10}$  and therefore can be used to calibrate the magnetic field of the measurement Penning trap.

(E) Beam request

We request six 12h-shifts of stable  ${}^7\text{Li}$  at 30 keV from the off-line ion source.

Once the procedure is established and sufficient accuracy and transfer efficiency is achieved the request for  ${}^{11}\text{Li}$  would be 4 shifts. During the  ${}^{11}\text{Li}$  run, beam-time would be required from the off-line source for field calibrations. The  ${}^7\text{Li}$  beam would be needed for approximately 20 min each every 4 h.

***Publications in refereed journals refereed conference proceedings:****An electromagnetic ion trap for studies in nuclear beta decay***AIP Conf. Proc. 457 1999, 172**

D. Beck; M. Beck; G. Bollen; J. Deutsch; J. Dilling; T. Phalet; P. Schuurmans; R. Prieels; W. Quint; N. Severijns; B. Vereecke; S. Versyck

*Search for scalar contributions to the  $^{38m}\text{K}$  beta- nu correlation in a magneto-optic trap***AIP Conf. Proc. 457 1999, 148**

J. A. Behr; A. Gorelov; D. Melconian; M. Trinczek; P. Dubé; O. Häusser; U. Giesen; K. P. Jackson; T. Swanson; J. M. D'Auria; M. Dombisky; G. Ball; L. Buchmann; B. Jennings; J. Dilling; J. Schmid; J. Deutsch; W. P. Alford; D. Asgeirsson; W. Wong

*Mass measurements of radioactive isotopes using the ISOLTRAP spectrometer***World Scientific Conf.Proc. 'Electroweak Physics' 1999, 324-329**

J. Dilling; F. Herfurth; H.-J. Kluge; A. Kohl; E. Lamour; G. Marx; S. Schwarz; G. Bollen; A. Kellerbauer; R.B. Moore

*The SHIPTRAP project: A capture and storage facility at GSI for heavy radionuclides from SHIP***Hyperfine Interactions 127 2000, 491-496**

Dilling, J.; Ackermann, D.; Bernard, J.; Hessberger, F.P.; Hofmann, S.; Hornung, W.; Kluge, H.-J.; Lamour, E.; Maier, M.; Mann, R.; Marx, G.; Moore, R.B.; Münzenberg, G.; Quint, W.; Rodriguez, D.; Schädel, M.; Schönfelder, J.; Sikler, G.; Toader, C.; Vermeeren, L.; Weber, C.; Bollen, G.; Engels, O.; Habs, D.; Thirof, P.; Backe, H.; Dretzke, A.; Lauth, W.; Ludolphs, W.; Sewtz, M.

*Beta-neutrino correlation experiments on laser trapped  $^{38m}\text{K}$ ,  $^{37}\text{K}$* **Hyperfine Interactions 127 2000, 373-380**

Gorelov, A.; Behr, J.A.; Melconian, D.; Trinczek, M.; Dubé, P.; Häusser, O.; Giesen, U.; Jackson, K.P.; Swanson, T.; D'Auria, J.M.; Dombisky, M.; Ball, G.; Buchmann, L.; Jennings, B.; Dilling, J.; Schmid, J.; Ashery, D.; Deutsch, J.; Alford, W.P.; Asgeirsson, D.; Wong, W.; Lee, B.

*Mass Measurements of  $^{114-124,130}\text{Xe}$  with the ISOLTRAP Penning Trap Spectrometer***Hyperfine Interactions 132 2001, 329-333**

Dilling, J.; Audi, G.; Beck, D.; Bollen, G.; Herfurth, F.; Kellerbauer, A.; Kluge, H.-J.; Lunney, D.; Moore, R.B.; Scheidenberger, C.; Schwarz, S.; Sikler, G.; Szerypo, J

*Towards Shorter-Lived Nuclides in ISOLTRAP Mass Measurements***Hyperfine Interactions 132 2001, 307-312**

Herfurth, F.; Dilling, J.; Kellerbauer, A.; Audi, G.; Beck, D.; Bollen, G.; Henry, S.; Kluge, H.-J.; Lunney, D.; Moore, R.B.; Scheidenberger, C.; Schwarz, S.; Sikler, G.; Szerypo, J

*Status of the SHIPTRAP Project: A Capture and Storage Facility for Heavy Radionuclides from SHIP***Hyperfine Interactions 132 2001, 459-464**

Marx, G.; Ackermann, D.; Dilling, J.; Hessberger, F.P.; Hoffmann, S.; Kluge, H.-J.; Mann, R.; Münzenberg, G.; Qamhieh, Z.b; Quint, W.b; Rodriguez, D.b; Schädel, M.; Schönfelder, J.; Sikler, G.; Toader, C.; Weber, C.; Engels, O.; Habs, D.; Thirof, P.; Backe, H.; Dretzke, A.; Lauth, W.; Ludolphs, W.; Sewtz, M.

*Improvement of the Applicability, Efficiency, and Precision of the Penning Trap Mass Spectrometer ISOLTRAP*

**Hyperfine Interactions 132 2001, 507-511**

Kellerbauer, A.; Bollen, G.; Dilling, J.; Henry, S.; Herfurth, F.; Kluge, H.-J.; Lamour, E.; Lunney, D.; Moore, R.B.; Scheidenberger, C.; Schwarz, S.; Sikler, G.; Szerypo, J

*A linear radiofrequency ion trap for accumulation, bunching, and emittance improvement of radioactive ion beams*

**Nuclear Instruments and Methods in Physics Research A 469 (2) 2001, 254-275**

Herfurth, F.; Dilling, J.; Kellerbauer, A.; Bollen, G.; Henry, S.; Kluge, H.-J.; Lamour, E.; Lunney, D.; Moore, R.B.; Scheidenberger, C.; Schwarz, S.; Sikler, G.; Szerypo, J

*A Physics Case for SHIPTRAP: Measuring the Masses of Transuranium Elements*

**Hyperfine Interactions 132 2001, 491-495**

Dilling, J.; Ackermann, D.; Heßberger, F. P.; Hofmann, S.; Kluge, H.-J.; Marx, G.; Münzenberg, G.; Schönfelder, J.; Sikler, G.; Sobiczewski, A.; Toader, C.; Weber, Chr.

*Stopping, Trapping and Cooling of Radioactive Fission Fragments in an Ion Catcher Device*

**Hyperfine Interactions 132 2001, 517-521**

Maier, M.; Boudreau, C.; Buchinger, F.; Clark, J.A.; Crawford, J.E.; Dilling, J.; Fukutani, H.; Gulick, S.; Lee, J.K.P.; Moore, R.B.; Savard, G.; Schwartz, J.; Sharma, K.S

*Mass Measurements on Short-Lived Nuclides with ISOLTRAP*

**Hyperfine Interactions 132 2001, 213-220**

Bollen, G.; Ames, F.; Audi, G.; Beck, D.; Dilling, J.; Engels, O.; Henry, S.; Herfurth, F.; Kellerbauer, A.; Kluge, H.-J.; Kohl, A.; Lamour, E.; Lunney, D.; Moore, R. B.; Oinonen, M.; Scheidenberger, C.; Schwarz, S.; Sikler, G.; Szerypo, J.; Weber, Chr.

*Accurate Mass Determination of Neutron-Deficient Nuclides Close to Z=82 with ISOLTRAP*

**Hyperfine Interactions 132 2001, 335-338**

Schwarz, S.; Ames, F.; Audi, G.; Beck, D.; Bollen, G.; Dilling, J.; Herfurth, F.; Kluge, H.-J.; Kellerbauer, A.; Kohl, A.; Lunney, D.; Moore, R.B.; Raimbault-Hartmann, H.; Scheidenberger, C.; Sikler, G.; Szerypo, J.

*HITRAP: A Facility for Experiments with Trapped Highly Charged Ions*

**Hyperfine Interactions 132 2001, 453-457**

Quint, W.; Dilling, J.; Djekic, S.; Häffner, H.; Hermanspahn, N.; Kluge, H.-J.; Marx, G.; Moore, R.c; Rodriguez, D.a; Schönfelder, J.; Sikler, G.; Valenzuela, T.; Verdú, J.; Weber, C.; Werth, G.

*Direct mass measurements of neutron-deficient xenon isotopes with the ISOLTRAP mass spectrometer*

**Nuclear Physics A 701 (1-4) 2002, 520-523**

Dilling, J.; Audi, G.; Beck, D.; Bollen, G.; Henry, S.; Herfurth, F.; Kellerbauer, A.; Kluge, H.-J.; Lunney, D.; Moore, R.B.; Scheidenberger, C.; Schwarz, S.; Sikler, G.; Szerypo, J.

*A linear radiofrequency quadrupole ion trap for the cooling and bunching of radioactive ion beams*

**Nuclear Physics A 701 (1-4) 2002, 565-569**

Kellerbauer, A.; Bollen, G.; Dilling, J.; Henry, S.; Herfurth, F.; Kluge, H.-J.; Lamour, E.; Moore, R.B.; Scheidenberger, C.; Schwarz, S.; Sikler, G.; Szerypo, J.

*Search for new physics in beta-neutrino correlations with the WITCH spectrometer*

**Nuclear Physics A 701 (1-4) 2002, 369-372**

Beck, D.; Ames, F.; Beck, M.; Bollen, G.; Delauré, B.; Deutsch, J.; Dilling, J. Forstner, O.; Phalet, T.; Prieels, R.; Quint, W.; Schmidt, P.; Schuurmans, P.; Severijns, N. Vereecke, B.; Versyck, S.

*Extension of Penning-trap mass measurements to very short-lived nuclides*

**Nuclear Physics A 701 (1-4) 2002, 516-519**

Herfurth, F.; Audi, G.; Beck, D.; Bollen, G.; Dilling, J.; Henry, S.; Kellerbauer, A.; Kluge, H.-J.; Kolhinen, V.; Lunney, D.; Moore, R.B.; Scheidenberger, C.; Schwarz, S.; Sikler, G.; Szerypo, J.

*SHIPTRAP—a capture and storage facility for heavy radionuclides at GSI*

**Nuclear Physics A 701 (1-4) 2002, 579-582**

Schönfelder, J.; Ackermann, D.; Backe, H.; Bollen, G.; Dilling, J.; Dretzke, A.; Engels, O.; Estermann, J.; Habs, D.; Hofmann, S.; Hessberger, F.P.; Kluge, H.-J.; Lauth, W.; Ludolphs, W.; Maier, M.; Marx, G.; Moore, R.B.; Quint, W.; Rodriguez, D.; Sewtz, M.; Sikler, G.; Toader, C.; Weber, Chr.

*Accurate masses of neutron-deficient nuclides close to  $Z=82$*

**Nuclear Physics A 701 (1-4) 2002, 533-545**

Schwarz, S.; Ames, F.; Audi, G.; Beck, D.; Bollen, G.; De Coster, C.; Dilling, J.; Engels, O.; Fossion, R.; Garcia Ramos, J.-E.; Henry, S.; Herfurth, F.; Heyde, K.; Kellerbauer, A.; Kluge, H.-J.; Kohl, A.; Lamour, E.; Lunney, D.; Martel, I.; Moore, R.B.; Oinonen, M.; Raimbault-Hartmann, H.; Scheidenberger, C.; Sikler, G.; Szerypo, J.; Weber, C.

*SHIPTRAP: A capture and storage facility on its way towards an RIB-facility*

**AIP Conf. Proc. 606 2002, 615-624**

G. Marx; J. Dilling; H.-J. Kluge; M. Mukherjee; W. Quint; S. Rahaman; D. Rodriguez; G. Sikler; M. Tarisien; C. Weber.

*Breakdown of the Isobaric Multiplet Mass Equation at  $A = 33$ ,  $T = 3/2$*

**Phys. Rev. Lett. 87 2002, 142501**

F. Herfurth; J. Dilling; A. Kellerbauer; G. Audi; D. Beck; G. Bollen; H.-J. Kluge; D. Lunney; R. B. Moore; C. Scheidenberger; S. Schwarz; G. Sikler; J. Szerypo.

*Novel Search for Heavy  $\nu$  Mixing from the beta Decay of  $K$  Confined in an Atom Trap*

**Phys. Rev. Lett. 90 2003, 012501**

M. Trinczek, A. Gorelov, D. Melconian, W. P. Alford, D. Asgeirsson, D. Ashery, J. A. Behr, P.G. Bricault, J.M. D'Auria, J. Deutsch, J. Dilling, M. Domsbky, P. Dube', S. Eaton, J. Fingler, U. Giesen, S. Gu, O. Ha'usser, K. P. Jackson, B. Lee, J. H. Schmid, T. J. Stocki, T. B. Swanson, and W. Wong

*SHIPTRAP is trapping: A capture and storage device on its way towards a RIB-facility*

**Hyperfine Interactions 146 (1-4) 245-251 2003**

Marx G., Dilling J., Kluge H.J., Mukherjee M., Quint W., Rahaman S., Rodriguez D., Sikler G., Tarisien M., Weber C.

*First on-line test of SHIPTRAP***Nuclear Instruments and Methods in Physics Research 204 2003 482-486**

Sikler G, Ackermann D, Attallah F, Beck D, Dilling J, Elisseev SA, Geissel H, Habs D, Habs D, Heinz S, Herfurth F, Hessberger F, Hofmann S, Kluge HJ, Kozhuharov C, Marx G, Mukherjee M, Neumayr J, Plass WR, Quint W, Rahaman S, Rodriguez D, Scheidenberger C, Tarisien M, Thirof P, Varentsov V, Weber C, Zhou Z

*The proposed TITAN facility at ISAC for very precise mass measurements on highly charged short-lived isotopes***Nuclear Instruments and Methods in Physics Research B 204 2003, 492-496**

J.Dilling; P. Bricault; M. Smith; H. -J. Kluge and the TITAN collaboration

*Mass of  $^{22}\text{Mg}$* **Phys. Rev. Lett. 93, 150801 (2004)**

M. Mukherjee, A. Kellerbauer, D. Beck, K. Blaum, G. Bollen, F. Carrel, P. Delahaye, J. Dilling, S. George, C. Gue'naut, F. Herfurth, A. Herlert, H.-J. Kluge, U. Koster, D. Lunney, S. Schwarz, L. Schweikhard, and C. Yazidjian

*Direct mass measurements of neutron-deficient xenon isotopes using the ISOLTRAP mass spectrometer***Euro. Physical Journal A 22, 163-171 (2004)**

J. Dilling, F. Herfurth, A. Kellerbauer, G. Audi, D. Beck, G. Bollen, H.-J. Kluge, R.B. Moore, C. Scheidenberger, S. Schwarz, G. Sikler, and the ISOLDE Collaboration

*Scalar Interaction Limits from the beta-nu Correlation of Trapped Radioactive Atoms***Phys. Rev. Lett. 94, 142501 (2005)**

A. Gorelov, D. Melconian, W. P. Alford, D. Ashery, G. Ball, J. A. Behr, P. G. Bricault, J. M. D'Auria, J. Deutsch, J. Dilling, M. Dombisky, P. Dube', J. Fingler, U. Giesen, F. Gluck, S. Gu, O. Hausser, K. P. Jackson, B. K. Jennings, M. R. Pearson, T. J. Stocki, T. B. Swanson, and M. Trinczek

*TITAN project status report and a proposal for a new cooling method of highly charged ions***Euro. Phys. Journal A 25 (2005) 1.53**

V. L. Ryjkov, L. Blomeley, M. Brodeur, P. Grothkopp, M. Smith, P. Bricault, F. Buchinger, J. Crawford, G. Gwinner, J. Lee, J. Vaz, G. Werth, J. Dilling and the TITAN Collaboration

*A high-current EBIT for charge-breeding of radionuclides for the TITAN spectrometer***Euro. Phys. Journal A 25 (2005) 1.63**

G. Sikler, J. R. Crespo López-Urrutia, J. Dilling, S. Epp, C. J. Osborne and J. Ullrich

*A high frequency MOSFET driver for the TITAN RFQ facility at TRIUMF***Proceedings of the IEEE International Pulsed Power Conference San Fransisco, Ca, USA. 2005**

M. J. Barnes, G. D. Wait, J. Dilling, J. V. Vaz, L. Blomeley O. Hadary, M. J. Smith.

*Nuclear Charge Radius of  $^{11}\text{Li}$ : Halo Neutron – Core Interactions***Submitted to Phys. Rev. Lett. and physics/0509265**

R. S'anchez, W. N'ortersh'ausser, G. Ewald, D. Albers J. Behr, P. Bricault, B.A. Bushaw, A. Dax, J. Dilling, M. Dombisky, G.W.F. Drake, S. G'otte, R. Kirchner, H.-J. Kluge, Th. K'uhl, J. Lassen, C.D.P. Levy, M. Pearson, E. Prime, V. Ryjkov, A. Wojtaszek, Z.-C. Yan, and C. Zimmermann

*Weak interaction symmetries with atom traps*

**Euro. Phys. Journal A 25 (2005 ) 1.685**

J. A. Behr, A. Gorelov, D. Melconian, M. Trinczek, W. P. Alford, D. Ashery, P. G. Bricault, L. Courneyea, J. M. D'Auria, J. Deutsch, J. Dilling, M. Dombisky, P. Dubé, F. Glück, S. Gryb, S. Gu, O. Häusser, K. P. Jackson, B. Lee, A. Mills, E. Paradis, M. Pearson, R. Pitcairn, E. Prime, D. Roberge and T. B. Swanson